

Global existence and well-posedness of 2D viscous shallow water system in Sobolev spaces with low regularity

Yanan Liu¹ Zhaoyang Yin²

Department of Mathematics, Sun Yat-sen University,
510275, Guangzhou, P. R. China

Abstract

In this paper we consider the Cauchy problem for 2D viscous shallow water system in $H^s(\mathbb{R}^2)$, $s > 1$. We first prove the local well-posedness of this problem by using the Littlewood-Paley theory, the Bony decomposition, and the theories of transport equations and transport diffusion equations. Then, we get the global existence of the system with small initial data in $H^s(\mathbb{R}^2)$, $s > 1$. Our obtained result improves the recent result in [12].

2010 Mathematics Subject Classification: 49K40, 42B25, 35A01, 35B44, 30H25.

Keywords: Viscous shallow water system; the Littlewood-Paley theory; the Bony Decomposition; Sobolev spaces; global existence

¹Email: *babyinarm@126.com*

²Email: *mcsyzy@mail.sysu.com.cn*

Contents

1	Introduction	2
2	Preliminaries	4
3	The local existence	10
3.1	Estimation for big norm	11
3.2	Convergence of small norm	15
3.3	Existence and uniqueness of local solutions	22
4	Global existence	23
	References	24

1 Introduction

We consider the following Cauchy problem for the following 2D viscous shallow water equations

$$(1.1) \quad \begin{cases} h(u_t + (u \cdot \nabla)u) - \nu \nabla \cdot (h \nabla u) + h \nabla h = 0, \\ h_t + \operatorname{div}(hu) = 0, \\ u|_{t=0} = u_0, \quad h|_{t=0} = h_0, \end{cases}$$

where $h(x, t)$ is the height of fluid surface, $u(x, t) = (u^1(x, t), u^2(x, t))$ is the horizontal velocity field, $x = (x_1, x_2) \in \mathbb{R}^2$, and $0 < \nu < 1$ is the viscous coefficient. For the initial data $h_0(x)$, we suppose that it is a small perturbation of some positive constant $\bar{h}_0$. In the paper, we mainly study the Cauchy problem (1.1) in Sobolev spaces $H^s(\mathbb{R}^2)$, $s > 1$. For the sake of convenience, we let the notation $B_{p,r}^s$ stand for $B_{p,r}^s(\mathbb{R}^2)$ in the

following text, and also let the notations L^p and H^s stand for $L^p(\mathbb{R}^2)$ and $H^s(\mathbb{R}^2)$, respectively.

Recently, Bresch et al. [3, 4] have systematically introduced the viscous shallow water equations. Bui in [5] proved the local existence and uniqueness of classical solutions to the Cauchy-Dirichlet problem for the shallow water equations with initial data in $C^{2+\alpha}$ by using Lagrangian coordinates and Hölder space estimates. Kloeden in [8] and Sundbye in [10] independently showed the global existence and uniqueness of classical solutions to the Cauchy-Dirichlet problem using Sobolev space estimates by following the energy method of Matsumura and Nishida [9]. Sundbye in [11] proved the existence and uniqueness of classical solutions to the Cauchy problem using the method of [9]. Wang and Xu in [12], obtained local solutions for any initial data and global solutions for small initial data $h_0 - \bar{h}_0, u_0 \in H^s$, $s > 2$. Haspot got global existence in time for small initial data $h_0, h_0 - \bar{h}_0 \in \dot{B}_{2,1}^0 \cap \dot{B}_{2,1}^1$ and $u_0 \in \dot{B}_{2,1}^0$ as a special case in [7], and Chen, Miao and Zhang in [6] to prove the local existence in time for general initial data and the global existence in time for small initial data where $h_0 - \bar{h}_0 \in \dot{B}_{2,1}^0 \cap \dot{B}_{2,1}^1$ and $u_0 \in \dot{B}_{2,1}^0$ with additional conditions that $h \geq h_0$ and h_0 is a strictly positive constant.

In the paper, we mainly use the Littlewood-Paley theory, the Bony decomposition and the Besov space theories for transport equations and transport-diffusion equations to obtain the local existence and uniqueness of solutions for any initial data in H^s , $s > 1$. And thus we can get the global existence of the system (1.1) with the small enough initial data in H^s , $s > 1$. The main result of this paper is as follows.

Theorem 1.1. *Let $u_0, h_0 - \bar{h}_0 \in H^s$, $s > 1$, $\|h_0 - \bar{h}_0\|_{H^s} < \bar{h}_0$. Then there exists a positive time T , a unique solution (u, h) of the Cauchy problem (1.1) such that*

$$u, h - \bar{h}_0 \in L^\infty([0, T]; H^s), \quad u \in L^2([0, T]; H^{s+1}).$$

Moreover, there exists a constant η , such that if $\|u_0\|_{H^s} + \|h_0\|_{H^s} \leq \eta$, then the time

T can be infinity.

2 Preliminaries

First of all, we transform the system (1.1). For a sake of convenience, we take $\bar{h}_0 = 1$.

Substituting h by $1 + h$ in (1.1), we have

$$(2.1) \quad \begin{cases} u_t + (u \cdot \nabla)u - \nu \Delta u - \nu \nabla(\ln(1 + h)) \nabla u + \nabla h = 0, \\ h_t + \operatorname{div} u + \operatorname{div}(hu) = 0, \\ u|_{t=0} = u_0, \quad h|_{t=0} = h_0, \end{cases}$$

$$x \in \mathbb{R}^2, t \in [0, \infty).$$

here $h_0 \in H^s$, $\|h_0\|_{H^s} \leq \frac{1}{8C_0}$, and C_0 will be determined below.

Next we introduce the well-known Littlewood-Paley decomposition briefly.

Proposition 2.1. [2] *Littlewood-Paley decomposition:*

Let $\mathcal{B} = \{\xi \in \mathbb{R}^2, |\xi| \leq \frac{4}{3}\}$ be a ball, and let $\mathcal{C} = \{\xi \in \mathbb{R}^2, \frac{3}{4} \leq |\xi| \leq \frac{8}{3}\}$ be an annulus. There exist two radial functions χ and φ valued in the interval $[0, 1]$, belonging respectively to $\mathcal{D}(\mathcal{B})$ and $\mathcal{D}(\mathcal{C})$, such that

$$(2.2) \quad \forall \xi \in \mathbb{R}^2, \quad \chi(\xi) + \sum_{j \geq 0} \varphi(2^{-j}\xi) = 1,$$

$$(2.3) \quad \forall \xi \in \mathbb{R}^2 \setminus \{0\}, \quad \sum_{j \in \mathbb{Z}} \varphi(2^{-j}\xi) = 1,$$

$$(2.4) \quad |j - j'| \geq 2 \Rightarrow \operatorname{Supp} \varphi(2^j \cdot) \cap \operatorname{Supp} \varphi(2^{j'} \cdot) = \emptyset,$$

$$(2.5) \quad j \geq 2 \Rightarrow \operatorname{Supp} \chi \cap \operatorname{Supp} \varphi(2^j \cdot) = \emptyset,$$

the set $\tilde{\mathcal{C}} \stackrel{\text{def}}{=} B(0, 2/3) + \mathcal{C}$ is an annulus, and we have

$$(2.6) \quad |j - j'| \geq 5 \Rightarrow 2^j \tilde{\mathcal{C}} \cap 2^{j'} \mathcal{C} = \emptyset.$$

Further, we have

$$(2.7) \quad \forall \xi \in \mathbb{R}^2, \quad \frac{1}{2} \leq \chi^2(\xi) + \sum_{j \geq 0} \varphi^2(2^{-j}\xi) \leq 1,$$

$$(2.8) \quad \forall \xi \in \mathbb{R}^2 \setminus \{0\}, \quad \frac{1}{2} \leq \sum_{j \in \mathbb{Z}} \varphi^2(2^{-j}\xi) \leq 1.$$

Now we can define the nonhomogeneous dyadic blocks Δ_j and the nonhomogeneous low-frequency cut-off operator S_j as follows:

$$\Delta_j u = 0, \text{ if } j \leq -2, \quad \Delta_{-1} u = \chi(D)u = \int_{\mathbb{R}^2} \tilde{h}(y)u(x-y)dy,$$

$$\Delta_j u = \varphi(2^{-j}D)u = 2^{jd} \int_{\mathbb{R}^2} h(2^j y)u(x-y)dy \quad \text{if } j \geq 0,$$

and

$$S_j u = \sum_{j' \leq j-1} \Delta_{j'} u,$$

where $h = \mathcal{F}^{-1}\varphi$ and $\tilde{h} = \mathcal{F}^{-1}\chi$.

Next we define the Besov spaces:

Definition 2.2. [2] Let $s \in \mathbb{R}$ and $(p, r) \in [1, \infty]^2$. The nonhomogeneous Besov space $B_{p,r}^s$ consists of all tempered distribution u such that:

$$\left(\sum_{j \geq -1} (2^{js} \|\Delta_j u\|_{L^p}) \right)_{\ell^r} < \infty,$$

and naturally the Besov norm is defined as follows

$$\|u\|_{B_{p,r}^s} = \left(\sum_{j \geq -1} (2^{js} \|\Delta_j u\|_{L^p}) \right)_{\ell^r}.$$

Particularly, when $p, r = 2$, the Besov space is coincident with the Sobolev space, i.e:

$$B_{2,2}^s = H^s.$$

Definition 2.3. [2] *The Bony decomposition: The nonhomogeneous paraproduct of v by u is defined by*

$$T_u v = \sum_j S_{j-1} u \Delta_j v.$$

The nonhomogeneous remainder of u by v is defined by

$$R(u, v) = \sum_{|k-j| \leq 1} \Delta_k u \Delta_j v.$$

The operators T and R are bilinear, and we have the following Bony decomposition

$$uv = T_v u + T_u v + R(u, v).$$

Lemma 2.4. [2] *Bernstein-Type inequalities:*

Let $\mathcal{C}$ be an annulus and $\mathcal{B}$ a ball. A constant C exists such that for any nonnegative integer k , any couple (p, q) in $[1, \infty]^2$ with $q \geq p \geq 1$, and any function u of L^p , we have

$$\text{Supp } \hat{u} \subset \lambda \mathcal{B} \Rightarrow \sup_{|\alpha|=k} \|\partial^\alpha u\|_{L^q} \leq C^{k+1} \lambda^{k+d(\frac{1}{p}-\frac{1}{q})} \|u\|_{L^p},$$

$$\text{Supp } \hat{u} \subset \lambda \mathcal{C} \Rightarrow C^{-k-1} \lambda^k \|u\|_{L^p} \leq \sup_{|\alpha|=k} \|\partial^\alpha u\|_{L^q} \leq C^{k+1} \lambda^k \|u\|_{L^p}.$$

Then we give some properties of the Besov spaces which will be used in this paper.

Lemma 2.5. [2] *Let $1 \leq p_1 \leq p_2 \leq \infty$ and $1 \leq r_1 \leq r_2 \leq \infty$. Then for any $s \in \mathbb{R}$, the space B_{p_1, r_1}^s is continuously embedded in $B_{p_2, r_2}^{s-d(\frac{1}{p_1}-\frac{1}{p_2})}$. Obviously, we also have that the space $B_{p, r}^{s_2}$ is continuously embedded in $B_{p, r}^{s_1}$ and $B_{p, \infty}^{s_2}$ is continuously embedded in $B_{p, 1}^{s_1}$ if $s_1 < s_2$.*

Lemma 2.6. [2] *If s_1 and s_2 are real numbers such that $s_1 < s_2$, $\theta \in (0, 1)$, and $p, r \in [1, \infty]$, then we have*

$$\|u\|_{B_{p, r}^{\theta s_1 + (1-\theta)s_2}} \leq \|u\|_{B_{p, r}^{s_1}}^\theta \|u\|_{B_{p, r}^{s_2}}^{1-\theta}.$$

Lemma 2.7. [2] If $u \in B_{p,r}^s$, then $\nabla u \in B_{p,r}^{s-1}$, and we have

$$\|\nabla u\|_{B_{p,r}^{s-1}} \leq C \|u\|_{B_{p,r}^s}.$$

Lemma 2.8. [2] The set $B_{p,r}^s$ is a Banach space. and satisfies the Fatou property, namely, if $(u_n)_{n \in \mathbb{N}}$ is a bounded sequence of $B_{p,r}^s$. Then an element u of $B_{p,r}^s$ and a subsequence $u_{\psi(n)}$ exist such that:

$$\lim_{n \rightarrow \infty} u_{\psi(n)} = u \text{ in } \mathcal{S}', \quad \|u\|_{B_{p,r}^s} \leq C \liminf_{n \rightarrow \infty} \|u_{\psi(n)}\|_{B_{p,r}^s}.$$

Lemma 2.9. [2] If $s > \frac{d}{p}$ or $s = \frac{d}{p}$, $r = 1$, then the $B_{p,r}^s$ space is continuously embedded in L^∞ , i.e

$$\|u\|_{L^\infty} \leq C_{s,p} \|u\|_{B_{p,r}^s}.$$

Lemma 2.10. [2] Let f be a smooth function, $f(0) = 0$, $s > 0$, $(p, r) \in [1, \infty]^2$. If $u \in B_{p,r}^s \cap L^\infty$, then so does $f \circ u$, and we have

$$\|f \circ u\|_{B_{p,r}^s} \leq C(s, f', \|u\|_{L^\infty}) \|u\|_{B_{p,r}^s}.$$

Lemma 2.11. [2] A constant C exists which satisfies the following inequalities for any couple of real numbers (s, t) with t negative and any (p, r_1, r_2) in $[1, \infty]^3$:

$$\|T\|_{\mathcal{L}(L^\infty \times B_{p,r}^s; B_{p,r}^s)} \leq C^{|s|+1},$$

$$\|T\|_{\mathcal{L}(B_{\infty,r_1}^t \times B_{p,r_2}^s; B_{p,r}^{s+t})} \leq \frac{C^{|s+t|+1}}{-t} \quad \text{with} \quad \frac{1}{r} \stackrel{\text{def}}{=} \min\left\{1, \frac{1}{r_1} + \frac{1}{r_2}\right\}.$$

Lemma 2.12. [2] A constant C exists which satisfies the following inequalities. Let (s_1, s_2) be in $\mathbb{R}^2$ and (p_1, p_2, r_1, r_2) be in $[1, \infty]^4$. Assume that

$$\frac{1}{p} \stackrel{\text{def}}{=} \frac{1}{p_1} + \frac{1}{p_2} \leq 1 \quad \text{and} \quad \frac{1}{r} \stackrel{\text{def}}{=} \frac{1}{r_1} + \frac{1}{r_2} \leq 1.$$

If $s_1 + s_2 > 0$, then we have, for any (u, v) in $B_{p_1,r_1}^{s_1} \times B_{p_2,r_2}^{s_2}$,

$$\|R(u, v)\|_{B_{p,r}^{s_1+s_2}} \leq \frac{C^{|s_1+s_2|+1}}{s_1 + s_2} \|u\|_{B_{p_1,r_1}^{s_1}} \|v\|_{B_{p_2,r_2}^{s_2}}.$$

If $r = 1$ and $s_1 + s_2 = 0$, then we have, for any (u, v) in $B_{p_1, r_1}^{s_1} \times B_{p_2, r_2}^{s_2}$,

$$\|R(u, v)\|_{B_{p, \infty}^0} \leq C^{|s_1 + s_2| + 1} \|u\|_{B_{p_1, r_1}^{s_1}} \|v\|_{B_{p_2, r_2}^{s_2}}.$$

Lemma 2.13. [2] For any $s > 0$ and $(p, r) \in [1, \infty]^2$, the space $B_{p, r}^s \cap L^\infty$ is an algebra, and a constant exists such that:

$$\|uv\|_{B_{p, r}^s} \leq \frac{C^{s+1}}{s} \left(\|u\|_{L^\infty} \|v\|_{B_{p, r}^s} + \|v\|_{L^\infty} \|u\|_{B_{p, r}^s} \right).$$

Moreover, if $s > \frac{d}{p}$ or $s = \frac{d}{p}, r = 1$, we have

$$\|uv\|_{B_{p, r}^s} \leq \frac{C^{s+1}}{s} \|u\|_{B_{p, r}^s} \|v\|_{B_{p, r}^s}.$$

For the transport equations

$$(2.9) \quad \begin{cases} \partial_t f + v \cdot \nabla f = g, \\ f|_{t=0} = f_0, \end{cases}$$

we have

Lemma 2.14. [2] Let $1 \leq p \leq p_1 \leq \infty$, $1 \leq r \leq \infty$. Assume that

$$(2.10) \quad s \geq -d \min \left(\frac{1}{p_1}, \frac{1}{p'} \right) \quad \text{or} \quad s \geq -1 - d \min \left(\frac{1}{p_1}, \frac{1}{p'} \right) \quad \text{if } \operatorname{div} v = 0$$

with strict inequality if $r < \infty$.

There exists a constant C , depending only on d, p, p_1, r and s , such that for all solutions $f \in L^\infty([0, T]; B_{p, r}^s)$ of (2.9), initial data f_0 in $B_{p, r}^s$, and g in $L^1([0, T]; B_{p, r}^s)$, we have, for a.e. $t \in [0, T]$,

$$\|f\|_{\tilde{L}_t^\infty(B_{p, r}^s)} \leq \left(\|f_0\|_{B_{p, r}^s} + \int_0^t \exp(-CV_{p_1}(t')) \|g(t')\|_{B_{p, r}^s} dt' \right) \exp(CV_{p_1}(t))$$

with, if the inequality is strict in (2.10),

$$(2.11) \quad V_{p_1}'(t) = \begin{cases} \|\nabla v(t)\|_{B_{p_1, r}^{s-1}}, & \text{if } s > 1 + \frac{d}{p_1} \text{ or } s = 1 + \frac{d}{p_1}, r = 1, \\ \|\nabla v(t)\|_{B_{p_1, \infty}^{\frac{d}{p_1}} \cap L^\infty}, & \text{if } s < 1 + \frac{d}{p_1}, \end{cases}$$

and if the equality holds in (2.10) and $r = \infty$,

$$V'_{p_1} = \|\nabla v(t)\|_{B_{p_1,1}^{\frac{d}{p_1}}}.$$

If $f = v$, then for all $s > 0$ ($s > -1$, if $\operatorname{div} u = 0$), the estimate holds with

$$V'_{p_1}(t) = \|\nabla u\|_{L^\infty},$$

where $\|u\|_{\tilde{L}_T^\rho(B_{p,r}^s)}$ is defined in Lemma 2.15.

For the transport diffusion equations

$$(2.12) \quad \begin{cases} \partial_t f + v \cdot \nabla f - \nu \Delta f = g, \\ f|_{t=0} = f_0, \end{cases}$$

we have the following lemma.

Lemma 2.15. [2] *Let $1 \leq p_1 \leq p \leq \infty$, $1 \leq r \leq \infty$, $s \in \mathbb{R}$ satisfy (2.10), and let V_{p_1} be defined as in Lemma 2.8.*

There exists a constant C which depends only on d, r, s and $s - 1 - \frac{d}{p_1}$ and is such that for any smooth solution of (11) and $1 \leq \rho_1 \leq \rho \leq \infty$, we have

$$\begin{aligned} \nu^{\frac{1}{\rho}} \|f\|_{\tilde{L}_T^\rho(B_{p,r}^{s+\frac{2}{\rho}})} &\leq C e^{C(1+\nu T)^{\frac{1}{\rho}} V_{p_1}(T)} \left((1 + \nu T)^{\frac{1}{\rho}} \|f_0\|_{B_{p,r}^s} \right. \\ &\quad \left. + (1 + \nu T)^{1+\frac{1}{\rho}-\frac{1}{\rho_1}} \nu^{\frac{1}{\rho_1}-1} \|g\|_{\tilde{L}_T^{\rho_1}(B_{p,r}^{s-2+\frac{2}{\rho_1}})} \right). \end{aligned}$$

For the space $\tilde{L}_T^\rho(B_{p,r}^s)$, we have the following properties:

Lemma 2.16. [2] *For all $T > 0$, $s \in \mathbb{R}$, and $1 \leq r, \rho \leq \infty$, we set*

$$\|u\|_{\tilde{L}_T^\rho(B_{p,r}^s)} \stackrel{\text{def}}{=} \|2^{js} \|\Delta_j u\|_{L_T^\rho(L^p)}\|_{l^r(\mathbb{Z})}.$$

We can then define the space $\tilde{L}_T^\rho(B_{p,r}^s)$ as the set of tempered distributions u over $(0, T) \times \mathbb{R}^d$ such that $\|u\|_{\tilde{L}_T^\rho(B_{p,r}^s)} \leq \infty$. By the Minkowski inequality, we have

$$\|u\|_{\tilde{L}_T^\rho(B_{p,r}^s)} \leq \|u\|_{L_T^\rho(B_{p,r}^s)} \quad \text{if } r \geq \rho,$$

$$\|u\|_{L_T^\rho(B_{p,r}^s)} \leq \|u\|_{\tilde{L}_T^\rho(B_{p,r}^s)} \quad \text{if } r \leq \rho.$$

The general principle is that all the properties of continuity for the product, composition, remainder, and paraproduct remain true in those spaces.

Moreover when $s > 0$, $1 \leq p \leq \infty$, $1 \leq \rho, \rho_1, \rho_2, \rho_3, \rho_4 \leq \infty$, and

$$\frac{1}{\rho} = \frac{1}{\rho_1} + \frac{1}{\rho_2} = \frac{1}{\rho_3} + \frac{1}{\rho_4},$$

we have

$$\|uv\|_{\tilde{L}_T^\rho(B_{p,r}^s)} \leq C \left(\|u\|_{\tilde{L}_T^{\rho_1}(L^\infty)} \|v\|_{\tilde{L}_T^{\rho_2}(B_{p,r}^s)} + \|v\|_{\tilde{L}_T^{\rho_3}(L^\infty)} \|u\|_{\tilde{L}_T^{\rho_4}(B_{p,r}^s)} \right).$$

Remark 2.17. For the sake of convenience, for the fixed s, p, r , we let $C_0(\geq 1)$ be the maximal constant of Lemmas 2.5-2.16.

3 The local existence

First of all, for the sake of convenience, we only consider the case of $s < 2$, the case of $s > 2$ was proved by Wang and Xu in [12], the case of $s = 2$ is similar to $s < 2$ but more easier. In order to study the local existence of solution, we define the function set $(u, h) \in \chi([0, T], s, E_1, E_2)$, if

$$u \in \tilde{L}^\infty([0, T]; H^s) \cap \tilde{L}^2([0, T]; H^{s+1}) \cap \tilde{L}^{\frac{4}{3-s}}([0, T]; H^{\frac{s+3}{2}}),$$

$$h \in \tilde{L}^\infty([0, T]; H^s),$$

and

$$\|u\|_{\tilde{L}^\infty([0, T]; H^s)} \leq E_1, \quad \|u\|_{\tilde{L}^2([0, T]; H^{s+1})} \leq E_1, \quad \|u\|_{\tilde{L}^{\frac{4}{3-s}}([0, T]; H^{\frac{s+3}{2}})} \leq E_1,$$

$$\|h\|_{\tilde{L}^\infty([0, T]; H^s)} \leq E_2,$$

where

$$E_1 = 8\nu^{-1}C_0\|u_0\|_{H^s}, \quad E_2 = 4\|h_0\|_{H^s}.$$

Next, we will prove Theorem 1.1 by the method of successive approximations. Let us define the sequence (u_n, h_n) by the following linear system:

$$(3.1) \quad \begin{cases} (u_1, h_1) = S_2(u_0, h_0), \\ \partial_t u_{n+1} + (u_n \cdot \nabla) u_{n+1} - \nu \Delta u_n = \frac{\nu}{1+h_n} \nabla h_n \nabla u_n + \nabla h_n, \\ \partial_t h_{n+1} + (u_n \cdot \nabla) h_{n+1} = -\operatorname{div} u_n - h \operatorname{div} u_n, \\ (u_{n+1}, h_{n+1})|_{t=0} = S_{n+2}(u_0, h_0). \end{cases}$$

Since S_q are smooth operators, the initial data $S_{n+2}(u_0, h_0)$ are smooth functions. If $(u_n, h_n) \in \chi([0, T], s, E_1, E_2)$ are smooth, then we have that for any $t \in [0, T]$,

$$\|h_n\|_{L^\infty} \leq C_0 \|h_n\|_{H^s} \leq C_0 E_2 = 4C_0 \|h_0\|_{H^s} \leq \frac{4C_0}{8C_0} = \frac{1}{2}.$$

Thus $\frac{\nu}{1+h_n} \nabla h_n \nabla u_n + \nabla h_n$ and $-\operatorname{div} u_n - h \operatorname{div} u_n$ are also smooth functions. Note that the first equation in (3.1) is a transport diffusion equation for u_{n+1} , and the second equation is a transport equation for h_{n+1} . Then the local existence of the smooth function for the Cauchy problem (3.1) is obvious.

We split the proof of Theorem 1.1 into two steps: Estimation for big norms and Convergence for small norms.

3.1. Estimation for big norm

In this subsection, we want to prove the following proposition.

Proposition 3.1. *Suppose that $(u_0, h_0) \in H^s \times H^s$, $s \in (1, 2)$ and $\|h_0\|_{B_{p,r}^s} \leq \frac{1}{8C_0}$, then there exists a positive time T , such that for any $n \in N$, $(u_n, h_n) \in \chi([0, T_1], s, E_1, E_2)$.*

Proof: Let $T \in (0, 1]$ satisfy firstly

$$e^{2C_0^2 E_1 T} \leq 2, \quad (1 + \nu T) \leq 2, \quad T \leq \frac{9}{64}.$$

Then we prove the proposition by induction. Firstly letting $(u_1, h_1) = S_2(u_0, h_0)$, thus we have

$$\|u_1\|_{\tilde{L}_T^\infty(H^s)} \leq \|u_0\|_{H^s} \leq E_1, \quad \|h_1\|_{\tilde{L}_T^\infty(H^s)} \leq \|h_0\|_{H^s} \leq E_2.$$

By Proposition 2.1, we also have

$$\|u_1\|_{L_T^2(H^{s+1})} \leq T^{\frac{1}{2}} \|S_2 u_0\|_{L^\infty(H^{s+1})} \leq \frac{32}{3} T^{\frac{1}{2}} \|S_2 u_0\|_{H^s} \leq E_1.$$

We suppose now that

$$\|u_n\|_{\tilde{L}_T^\infty(H^s)} \leq E_1, \quad \|u_1\|_{L_T^2(H^{s+1})} \leq E_1, \quad \|h_n\|_{\tilde{L}_T^\infty(H^s)} \leq E_2.$$

Then for u_{n+1} , in the view of Lemma 2.15, taking a positive real number ρ such that $\frac{2}{\rho} < s - 1$, then for all $t \leq T$, we have

$$\begin{aligned} (3.2) \quad & \|u_{n+1}\|_{\tilde{L}_t^\infty(H^s)} \leq C_0 e^{C_0 \int_0^t \|\nabla u_n(t')\|_{H^1 \cap L^\infty} dt'} \left(\|S_{n+2} u_0\|_{H^s} + (1 + \nu T)^{1 - \frac{1}{\rho}} \nu^{\frac{1}{\rho} - 1} \times \right. \\ & \left. \|\nu \nabla(\ln(1 + h_n)) \nabla u_n - \nabla h_n\|_{\tilde{L}_t^\rho(H^{s-2+\frac{2}{\rho}})} \right) \\ & \leq 2C_0 \left(\frac{E_1}{4C_0} + 2\nu^{\frac{1}{\rho} - 1} \|\nabla h_n\|_{\tilde{L}_t^\rho(H^{s-2+\frac{2}{\rho}})} + 2\nu^{\frac{1}{\rho}} \|\nabla(\ln(1 + h_n)) \nabla u_n\|_{\tilde{L}_t^\rho(H^{s-2+\frac{2}{\rho}})} \right) \\ & \leq \frac{E_1}{2} + CT^{\frac{1}{\rho}} \|h_n\|_{\tilde{L}_T^\infty(H^s)} + C \|\nabla(\ln(1 + h_n)) \nabla u_n\|_{\tilde{L}_t^\rho(H^{s-2+\frac{2}{\rho}})}. \end{aligned}$$

Now, we estimate $\|\nabla(\ln(1 + h_n)) \nabla u_n\|_{\tilde{L}_t^\rho(H^{s-2+\frac{2}{\rho}})}$. By Lemmas 2.11 and 2.12, we get

$$\begin{aligned} (3.3) \quad & \|\nabla(\ln(1 + h_n)) \nabla u_n\|_{\tilde{L}_t^\rho(H^{s-2+\frac{2}{\rho}})} \\ & \leq \|T_{\nabla(\ln(1+h_n))} \nabla u_n + T_{\nabla u_n} \nabla(\ln(1 + h_n)) + R(\nabla(\ln(1 + h_n)), \nabla u_n)\|_{L_t^\rho(H^{s-2+\frac{2}{\rho}+\varepsilon})} \\ & \leq CT^{\frac{1}{\rho}} (\|\nabla(\ln(1 + h_n))\|_{L_T^\infty(B_{\infty,\infty}^{s-2})} \|\nabla u_n\|_{L_T^\infty(H^{\frac{2}{\rho}+\varepsilon})} + \|\nabla u_n\|_{L_T^\infty(B_{\infty,\infty}^{s-2})} \|\nabla(\ln(1 + h_n))\|_{L_T^\infty(H^{\frac{2}{\rho}+\varepsilon})}) \\ & \quad + CT^{\frac{1}{\rho}} \|R(\nabla(\ln(1 + h_n)), \nabla u_n)\|_{L_T^\infty(B_{1,2}^{s-1+\frac{2}{\rho}+\varepsilon})} \\ & \leq CT^{\frac{1}{\rho}} \|u_n\|_{L_T^\infty(H^s)} \|h_n\|_{L_T^\infty(H^s)} + CT^{\frac{1}{\rho}} \|\nabla(\ln(1 + h_n))\|_{L_T^\infty(H^{s-1})} \|\nabla u_n\|_{L_T^\infty(H^{\frac{2}{\rho}+\varepsilon})} \\ & \leq CT^{\frac{1}{\rho}} \|u_n\|_{L_T^\infty(H^s)} \|h_n\|_{L_T^\infty(H^s)}, \end{aligned}$$

where ε is a small enough positive real number, and it doesn't affect the direction of the inequality:

$$\text{if } 1 + \frac{2}{\rho} < s, \text{ then } 1 + \frac{2}{\rho} + \varepsilon < s.$$

Thus combining with (3.2), we get

$$\begin{aligned} (3.4) \quad \|u_{n+1}\|_{\tilde{L}_t^\infty(H^s)} &\leq \frac{E_1}{2} + CT^{\frac{1}{\rho}} \|h_n\|_{\tilde{L}_T^\infty(H^s)} + CT^{\frac{1}{\rho}} \|u_n\|_{L_T^\infty(H^s)} \|h_n\|_{L_T^\infty(H^s)} \\ &\leq \frac{E_1}{2} + CT^{\frac{1}{\rho}} E_2 + CT^{\frac{1}{\rho}} E_1 E_2. \end{aligned}$$

Remark 3.2. In the calculation of (3.3), if $s = 2$, we can estimate

$$\|T_{\nabla(\ln(1+h_n))} \nabla u_n\|_{L_t^\rho(H^{s-2+\frac{2}{\rho}+\varepsilon})}$$

as follows:

$$\begin{aligned} (3.5) \quad &\|T_{\nabla(\ln(1+h_n))} \nabla u_n\|_{L_t^\rho(H^{s-2+\frac{2}{\rho}+\varepsilon})} \\ &\leq CT^{\frac{1}{\rho}} \|\nabla(\ln(1+h_n))\|_{L_T^\infty(B_{\infty,\infty}^{s-2-\varepsilon})} \|\nabla u_n\|_{L_T^\infty(H^{\frac{2}{\rho}+2\varepsilon})} \\ &\leq CT^{\frac{1}{\rho}} \|h_n\|_{L_T^\infty(H^s)} \|u_n\|_{L_T^\infty(H^s)}. \end{aligned}$$

Using the same method, we deal with the similar case below as well.

For h_{n+1} , from Lemmas 2.14, we obtain

$$\begin{aligned} (3.6) \quad &\|h_{n+1}\|_{\tilde{L}_t^\infty(H^s)} \leq e^{C_0 \int_0^t \|\nabla u_n\|_{H^1 \cap L^\infty} dt'} \left(\|S_{n+2} h_0\|_{H^s} + \|\operatorname{div} u_n\|_{\tilde{L}_T^1(H^s)} + \|h_n \operatorname{div} u_n\|_{\tilde{L}_T^1(H^s)} \right) \\ &\leq 2\|h_0\|_{H^s} + C\|u_n\|_{\tilde{L}_T^1(H^{s+1})} + C\|h_n\|_{L_T^\infty(H^s)} \|\operatorname{div} u_n\|_{L_T^1(H^s)} \\ &\leq \frac{E_2}{2} + (C + C\|h_n\|_{L_T^\infty(H^s)}) T^{\frac{1}{2}} \|u_n\|_{L_T^2(H^{s+1})} \\ &\leq \frac{E_2}{2} + CT^{\frac{1}{2}} (1 + E_2) E_1. \end{aligned}$$

Then by Lemma 2.15 again, we have

(3.7)

$$\begin{aligned}
\|u_{n+1}\|_{L_t^2(H^{s+1})} &\leq \nu^{-\frac{1}{2}} C_0 e^{C_0(1+\nu T)^{\frac{1}{2}} \int_0^T \|\nabla u_n\|_{H^1 \cap L^\infty} dt} \left((1+\nu T)^{\frac{1}{2}} \|S_{n+1} u_0\|_{H^s} + \right. \\
&\quad \left. (1+\nu T) \nu^{-\frac{1}{2}} \|\nu \nabla(\ln(1+h_n)) \nabla u_n - \nabla h_n\|_{L_T^2(H^{s-1})} \right) \\
&\leq 2C_0 \nu^{-\frac{1}{2}} (2\|u_0\|_{H^s} + 2\nu^{-\frac{1}{2}} C_0 \|h_n\|_{L_T^2(H^s)} + 2\nu^{-\frac{1}{2}} \|\nabla(\ln(1+h_n)) \nabla u_n\|_{L_T^2(H^{s-1})}) \\
&\leq 4C_0 \nu^{-\frac{1}{2}} \|u_0\|_{H^s} + CT^{\frac{1}{2}} \|h_n\|_{L_T^\infty(H^s)} + C \|\nabla(\ln(1+h_n)) \nabla u_n\|_{L_T^2(H^{s-1})}.
\end{aligned}$$

By Lemmas 2.11-2.12, we have

(3.8)

$$\begin{aligned}
\|\nabla(\ln(1+h_n)) \nabla u_n\|_{L_T^2(H^{s-1})} &\leq C \|\nabla(\ln(1+h_n))\|_{L_T^\infty(B_{\infty,\infty}^{s-2})} \|\nabla u_n\|_{L_T^2(H^1)} + \\
C \|\nabla u_n\|_{L_T^2(L^\infty)} \|\nabla(\ln(1+h_n))\|_{L_T^\infty(H^{s-1})} &+ C \|\nabla(\ln(1+h_n))\|_{L_T^\infty(B_{\infty,\infty}^{s-2})} \|\nabla u_n\|_{L_T^2(H^1)} \\
&\leq C \|h_n\|_{L_T^\infty(H^s)} \|u_n\|_{L_T^2(H^{\frac{s+3}{2}})} \\
&\leq CT^{\frac{s-1}{4}} \|h_n\|_{L_T^\infty(H^s)} \|u_n\|_{L_T^{\frac{4}{3-s}}(H^{\frac{s+3}{2}})}.
\end{aligned}$$

By Lemma 2.6, and taking $\theta = \frac{s-1}{2}$, we have

$$(3.9) \quad \|u_n\|_{L_T^{\frac{4}{3-s}}(H^{\frac{s+3}{2}})} \leq \|u_n\|_{L_T^\infty(H^s)}^{\frac{s-1}{2}} \|u_n\|_{L_T^2(H^{s+1})}^{\frac{3-s}{2}} \leq E_1.$$

Combining (3.7), (3.8) and (3.9), we can obtain that

$$\begin{aligned}
\|u_{n+1}\|_{L_t^2(H^{s+1})} &\leq 4C_0 \nu^{-\frac{1}{2}} \|u_0\|_{H^s} + CT^{\frac{1}{2}} \|h_n\|_{L_T^\infty(H^s)} \\
(3.10) \quad &+ CT^{\frac{s-1}{4}} \|h_n\|_{L_T^\infty(H^s)} \|u_n\|_{L_T^\infty(H^s)}^{\frac{s+1}{2}} \|u_n\|_{L_T^2(H^{s+1})}^{\frac{3-s}{2}} \\
&\leq \frac{E_1}{2} + CT^{\frac{1}{2}} E_2 + CT^{\frac{s-1}{4}} E_2 E_1,
\end{aligned}$$

where C in (3.2)-(3.10) only depends on C_0, ν . Thus we only take a suitable T , such that

$$(3.11) \quad \begin{cases} CT^{\frac{1}{\rho}} E_2 \leq \frac{E_1}{4}, & CT^{\frac{1}{\rho}} E_2 \leq \frac{1}{4}, \\ CT^{\frac{1}{2}} (1+E_2) E_1 \leq \frac{E_2}{2}, \\ CT^{\frac{1}{2}} E_2 \leq \frac{E_1}{4}, & CT^{\frac{s-1}{4}} E_2 \leq \frac{1}{4}. \end{cases}$$

This completes the proof of Proposition 3.1.

3.2. Convergence of small norm

Proposition 3.3. *Suppose that $(u_0, h_0) \in H^s \times H^s$, $s \in (1, 2)$, $\|h_0\|_{B_{p,r}^s} \leq \frac{1}{8C_0}$, then there exists a positive time $T_1(\leq T)$, such that (u_n, h_n) is a Cauchy sequence in $\chi([0, T_1], s-1, E_1, E_2)$.*

Proof: From the equations in (3.1), we have

$$(3.12) \quad \begin{cases} \partial_t(u_{n+1} - u_n) + (u_n \cdot \nabla)(u_{n+1} - u_n) - \nu \Delta(u_{n+1} - u_n) = \sum_{j=1}^5 F_j \\ \partial_t(h_{n+1} - h_n) + (u_n \cdot \nabla)(h_{n+1} - h_n) = \sum_{j=1}^4 J_j, \\ (u_{n+1} - u_n, h_{n+1} - h_n)|_{t=0} = \Delta_{n+1}(u_0, h_0), \end{cases}$$

where

$$(3.13) \quad \begin{aligned} \sum_{j=1}^5 F_j &= (u_n - u_{n-1}) \cdot \nabla u_n + \nabla(h_n - h_{n-1}) + \frac{\nu}{1+h_n} \nabla h_n \nabla(u_n - u_{n-1}) \\ &+ \frac{\nu}{1+h_n} \nabla u_{n-1} \nabla(h_n - h_{n-1}) + \nu \left(\frac{1}{1+h_n} - \frac{1}{1+h_{n-1}} \right) \nabla h_{n-1} \nabla u_{n-1}, \\ \sum_{j=1}^4 J_j &= (u_n - u_{n-1}) \cdot \nabla h_n + \operatorname{div}(u_n - u_{n-1}) + h_n \operatorname{div}(u_n - u_{n-1}) \\ &+ (h_n - h_{n-1}) \operatorname{div} u_{n-1}. \end{aligned}$$

Then we estimate the Sobolev norms of $u_{n+1} - u_n$ and $h_{n+1} - h_n$. For any $t \leq T_1 \leq T$, by Lemma 2.15, we have

$$(3.14) \quad \begin{aligned} \|u_{n+1} - u_n\|_{\tilde{L}_t^\infty(H^{s-1})} &\leq C_0 e^{\int_0^t \|\nabla u_n\|_{H^1 \cap L^\infty} dt'} \times \\ &\left(\|S_{n+2}u_0 - S_{n+1}u_0\|_{H^{s-1}} + \sum_{j=1}^5 \left(\frac{1+\nu t}{\nu} \right)^{1-\frac{1}{\rho_j}} \|F_j\|_{\tilde{L}_t^{\rho_j}(H^{s-3+\frac{2}{\rho_j}})} \right) \\ &\leq 2C_0 \left(\|\Delta_{n+1}u_0\|_{H^{s-1}} + 2\nu^{-1} \sum_{j=1}^5 \|F_j\|_{\tilde{L}_t^{\rho_j}(H^{s-3+\frac{2}{\rho_j}})} \right). \end{aligned}$$

Taking $\rho_1 = 2$, by Lemmas 2.11-2.12, we have

(3.15)

$$\begin{aligned}
& \|F_1\|_{\tilde{L}_t^2(H^{s-2})} = \|(u_n - u_{n-1}) \cdot \nabla u_n\|_{\tilde{L}_t^2(H^{s-2})} \\
& \leq C \|T_{u_n - u_{n-1}} \nabla u_n + T_{\nabla u_n} (u_n - u_{n-1}) + R(\nabla u_n, u_n - u_{n-1})\|_{\tilde{L}_t^2(H^{s-2})} \\
& \leq C \|u_n - u_{n-1}\|_{L_t^2(B_{\infty,\infty}^{s-2})} \|\nabla u_n\|_{L_t^\infty(H^0)} + \|\nabla u_n\|_{L_t^\infty(B_{\infty,\infty}^{-1})} \|u_n - u_{n-1}\|_{L_t^2(H^{s-1})} \\
& \quad + C \|\nabla u_n\|_{L_t^\infty(H^0)} \|u_n - u_{n-1}\|_{L_t^2(H^{s-1})} \\
& \leq CT_1^{\frac{1}{2}} \|u_n\|_{L_T^\infty(H^s)} \|u_n - u_{n-1}\|_{L_t^\infty(H^{s-1})} \\
& \leq CT_1^{\frac{1}{2}} E_1 \|u_n - u_{n-1}\|_{L_t^\infty(H^{s-1})}.
\end{aligned}$$

Taking $\rho_2 = 2$ as well, we have

$$(3.16) \quad \|F_2\|_{\tilde{L}_t^2(H^{s-2})} = \|\nabla(h_n - h_{n-1})\|_{\tilde{L}_t^2(H^{s-2})} \leq CT_1^{\frac{1}{2}} \|h_n - h_{n-1}\|_{\tilde{L}_t^\infty(H^{s-1})}.$$

Taking $\rho_3 = \frac{2}{s}$, by Lemmas 2.11-2.12, we get

(3.17)

$$\begin{aligned}
& \|F_3\|_{\tilde{L}_t^{\frac{2}{s}}(H^{s-3+s})} = \|\nu \nabla(\ln(1 + h_n)) \nabla(u_n - u_{n-1})\|_{L_t^{\frac{2}{s}}(H^{2s-3})} \\
& \leq C \|T_{\nabla(\ln(1+h_n))} \nabla(u_n - u_{n-1})\|_{L_t^{\frac{2}{s}}(H^{2s-3})} + C \|T_{\nabla u_n - u_{n-1}} \nabla(\ln(1 + h_n))\|_{L_t^{\frac{2}{s}}(H^{2s-3})} \\
& \quad + C \|R(\nabla(\ln(1 + h_n)), \nabla(u_n - u_{n-1}))\|_{L_t^{\frac{2}{s}}(H^{2s-3})} \\
& \leq C \|\nabla(\ln(1 + h_n))\|_{L_t^\infty(B_{\infty,\infty}^{s-2})} \|\nabla(u_n - u_{n-1})\|_{L_t^{\frac{2}{s}}(H^{s-1})} \\
& \quad + C \|\nabla(u_n - u_{n-1})\|_{L_t^{\frac{2}{s}}(B_{\infty,\infty}^{s-2})} \|\nabla(\ln(1 + h_n))\|_{L_t^\infty(H^{s-1})} \\
& \quad + C \|\nabla(u_n - u_{n-1})\|_{L_t^{\frac{2}{s}}(H^{s-1})} \|\nabla(\ln(1 + h_n))\|_{L_t^\infty(H^{s-1})} \\
& \leq C \|\ln(1 + h_n)\|_{L_t^\infty(H^s)} \|u_n - u_{n-1}\|_{L_t^{\frac{2}{s}}(H^s)} \\
& \leq CT_1^{\frac{s-1}{2}} \|h_n\|_{L_t^\infty(H^s)} \|u_n - u_{n-1}\|_{L_t^2(H^s)} \\
& \leq CT_1^{\frac{s-1}{2}} E_2 \|u_n - u_{n-1}\|_{L_t^2(H^s)}.
\end{aligned}$$

Then we deal with F_4 by the similar method. Taking $\rho_4 = 2$, we have

$$\begin{aligned}
(3.18) \quad & \|F_4\|_{L_t^2(H^{s-2})} = \|\nu \frac{\nabla(h_n - h_{n-1})}{1+h_n} \nabla u_{n-1}\|_{L_t^2(H^{s-2})} \\
& \leq C \|\nabla u_{n-1} \nabla(h_n - h_{n-1})\|_{L_t^2(H^{s-2})} + C \|\frac{h_n}{1+h_n} \nabla u_{n-1} \nabla(h_n - h_{n-1})\|_{L_t^2(H^{s-2})} \\
& = F_{41} + F_{42}.
\end{aligned}$$

By Lemmas 2.11-2.12, we have

$$\begin{aligned}
(3.19) \quad & F_{41} \leq C \|\nabla(h_n - h_{n-1})\|_{L_t^\infty(B_{\infty,\infty}^{s-3})} \|\nabla u_{n-1}\|_{L_T^2(H^1)} + \|\nabla u_{n-1}\|_{L_t^2(L^\infty)} \|\nabla(h_n - h_{n-1})\|_{L_t^\infty(H^{s-2})} \\
& + C \|\nabla u_{n-1}\|_{L_t^2(H^1)} \|\nabla(h_n - h_{n-1})\|_{L_t^\infty(H^{s-2})} \\
& \leq C \|\nabla u_n\|_{L_t^2(L^\infty \cap H^1)} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
& \leq C \|u_n - u_{n-1}\|_{L_t^2(H^{\frac{s+3}{2}})} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
& \leq CT_1^{\frac{s-1}{4}} \|u_{n-1}\|_{L_t^{\frac{4}{3-s}}(H^{\frac{s+3}{2}})} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
& \leq CT_1^{\frac{s-1}{4}} E_1 \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})}.
\end{aligned}$$

Similarly as F_{41} , using Lemmas 2.11-2.12 again, we have

$$\begin{aligned}
(3.20) \quad & F_{42} \leq C \|\frac{h_n}{1+h_n} \nabla u_{n-1}\|_{L_t^2(L^\infty \cap H^1)} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
& \leq C \|\frac{h_n}{1+h_n} \nabla u_{n-1}\|_{L_t^2(H^{\frac{s+1}{2}})} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
& \leq C \|\frac{h_n}{1+h_n}\|_{L_t^\infty(H^s)} \|\nabla u_{n-1}\|_{L_t^2(H^{\frac{s+1}{2}})} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
& \leq CT_1^{\frac{s-1}{4}} \|h_n\|_{L_t^\infty(H^s)} \|u_{n-1}\|_{L_t^{\frac{4}{3-s}}(H^{\frac{s+3}{2}})} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
& \leq CT_1^{\frac{s-1}{4}} E_1 E_2 \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})}.
\end{aligned}$$

For F_5 , taking $\rho_5 = 2$, then we have

$$\begin{aligned}
\|F_5\|_{L_t^2(H^{s-2})} &= \left\| \nu \left(\frac{1}{1+h_n} - \frac{1}{1+h_{n-1}} \right) \nabla h_{n-1} \nabla u_{n-1} \right\|_{L_t^2(H^{s-2})} \\
&= \nu \left\| (h_{n-1} - h_n) \frac{1}{1+h_n} \nabla(\ln(1+h_{n-1})) \nabla u_{n-1} \right\|_{L_t^2(H^{s-2})} \\
(3.21) \quad &\leq C \left\| (h_{n-1} - h_n) \nabla(\ln(1+h_{n-1})) \nabla u_{n-1} \right\|_{L_t^2(H^{s-2})} \\
&\quad + \left\| (h_{n-1} - h_n) \frac{h_n}{1+h_n} \nabla(\ln(1+h_{n-1})) \nabla u_{n-1} \right\|_{L_t^2(H^{s-2})} \\
&= F_{51} + F_{52}.
\end{aligned}$$

By Lemmas 2.11-2.12, we have

$$\begin{aligned}
F_{51} &\leq C \left\| T_{\nabla(\ln(1+h_{n-1})) \nabla u_{n-1}} (h_n - h_{n-1}) \right\|_{L_t^2(H^{s-2})} \\
&\quad + C \left\| T_{h_n - h_{n-1}} (\nabla(\ln(1+h_{n-1})) \nabla u_{n-1}) \right\|_{L_t^2(H^{s-2})} \\
&\quad + C \left\| R(\nabla(\ln(1+h_{n-1})) \nabla u_{n-1}, (h_n - h_{n-1})) \right\|_{L_t^2(H^{s-2})} \\
(3.22) \quad &\leq C \left\| \nabla(\ln(1+h_{n-1})) \nabla u_{n-1} \right\|_{L_t^2(B_{\infty,\infty}^{-1})} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
&\quad + C \|h_n - h_{n-1}\|_{L_t^\infty(B_{\infty,\infty}^{s-2})} \left\| \nabla(\ln(1+h_{n-1})) \nabla u_{n-1} \right\|_{L_t^2(H^0)} \\
&\quad + \left\| \nabla(\ln(1+h_{n-1})) \nabla u_{n-1} \right\|_{L_t^2(H^0)} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
&\leq C \left\| \nabla(\ln(1+h_{n-1})) \nabla u_{n-1} \right\|_{L_t^2(H^0)} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})}.
\end{aligned}$$

By Lemmas 2.11-2.12 again, we have

$$\begin{aligned}
&\left\| \nabla(\ln(1+h_{n-1})) \nabla u_{n-1} \right\|_{L_t^2(H^0)} \\
&\leq C \left\| \nabla(\ln(1+h_{n-1})) \right\|_{L_t^\infty(B_{\infty,\infty}^{s-2})} \left\| \nabla u_{n-1} \right\|_{L_t^2(H^{2-s})} \\
&\quad + C \left\| \nabla u_{n-1} \right\|_{L_t^2(B_{\infty,\infty}^{1-s})} \left\| \nabla(\ln(1+h_{n-1})) \right\|_{L_t^\infty(H^{s-1})} \\
(3.23) \quad &\quad + C \left\| \nabla(\ln(1+h_{n-1})) \right\|_{L_t^\infty(H^{s-1})} \left\| \nabla u_{n-1} \right\|_{L_t^2(H^{2-s})} \\
&\leq C \|h_{n-1}\|_{L_t^\infty(H^s)} \|u_{n-1}\|_{L_t^2(H^{3-s})} \\
&\leq C T_1^{\frac{s-1}{4}} \|h_{n-1}\|_{L_t^\infty(H^s)} \|u_{n-1}\|_{L_t^{\frac{4}{3-s}}(H^{\frac{s+3}{2}})}.
\end{aligned}$$

Combining (3.22) and (3.23), we have

$$\begin{aligned}
(3.24) \quad F_{51} &\leq CT_1^{\frac{s-1}{4}} \|h_{n-1}\|_{L_t^\infty(H^s)} \|u_{n-1}\|_{L_t^{\frac{4}{3-s}}(H^{\frac{s+3}{2}})} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
&\leq CT_1^{\frac{s-1}{4}} E_1 E_2 \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})}.
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
(3.25) \quad F_{52} &\leq C \left\| \frac{h_n}{1+h_n} \nabla(\ln(1+h_{n-1})) \nabla u_{n-1} \right\|_{L_t^2(H^0)} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
&\leq C \left\| \frac{h_n}{1+h_n} \right\|_{L_t^\infty(H^s)} \|\nabla(\ln(1+h_{n-1})) \nabla u_{n-1}\|_{L_t^2(H^0)} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
&\leq CT_1^{\frac{s-1}{4}} \|h_{n-1}\|_{L_t^\infty(H^s)}^2 \|u_{n-1}\|_{L_t^{\frac{4}{3-s}}(H^{\frac{s+3}{2}})} \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
&\leq CT_1^{\frac{s-1}{4}} E_1 E_2^2 \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})}.
\end{aligned}$$

Thus, we have

$$\begin{aligned}
(3.26) \quad &\|u_{n+1} - u_n\|_{\tilde{L}_t^\infty(H^{s-1})} \leq 2C_0 \|\Delta_{n+1} u_0\|_{H^{s-1}} + CT^{\frac{1}{2}} E_1 \|u_n - u_{n-1}\|_{L_t^\infty(H^{s-1})} \\
&+ CT_1^{\frac{1}{2}} \|h_n - h_{n-1}\|_{\tilde{L}_t^\infty(H^{s-1})} + CT_1^{\frac{s-1}{2}} E_2 \|u_n - u_{n-1}\|_{L_t^2(H^s)} \\
&+ CT_1^{\frac{s-1}{4}} E_1 \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} + CT_1^{\frac{s-1}{4}} E_1 E_2 \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
&+ CT_1^{\frac{s-1}{4}} E_1 E_2 \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} + CT^{\frac{s-1}{4}} E_1 E_2^2 \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
&\leq 2C_0 \|\Delta_{n+1} u_0\|_{H^{s-1}} + CT_1^{\frac{s-1}{4}} E_1 \|u_n - u_{n-1}\|_{L_t^\infty(H^{s-1})} + CT_1^{\frac{s-1}{4}} E_2 \|u_n - u_{n-1}\|_{L_t^2(H^s)} \\
&+ CT_1^{\frac{s-1}{4}} (1 + E_1 + E_1 E_2 + E_1 E_2^2) \|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})}.
\end{aligned}$$

For $h_{n+1} - h_n$, we have

$$\begin{aligned}
(3.27) \quad &\|h_{n+1} - h_n\|_{\tilde{L}_t^\infty(H^{s-1})} \leq e^{C_0 \int_0^t \|\nabla u_n\|_{H^1 \cap L^\infty}} \times (\|\Delta_{n+1} h_0\|_{H^{s-1}} + \|\sum_{j=1}^4 G_j\|_{\tilde{L}_t^1(H^{s-1})}) \\
&\leq 2(\|\Delta_{n+1} h_0\|_{H^{s-1}} + T_1^{\frac{1}{2}} \|\sum_{j=1}^4 J_j\|_{\tilde{L}_t^2(H^{s-1})}).
\end{aligned}$$

From Lemma 2.12, we have

(3.28)

$$\begin{aligned}
& \|J_1\|_{\tilde{L}_t^2(H^{s-1})} = \|(u_n - u_{n-1}) \cdot \nabla h_n\|_{\tilde{L}_t^2(H^{s-1})} \\
& \leq C\|u_n - u_{n-1}\|_{L_t^2(L^\infty)}\|\nabla h_n\|_{L_t^\infty(H^{s-1})} + C\|\nabla h_n\|_{L_t^\infty(B_{\infty,\infty}^{s-2})}\|u_n - u_{n-1}\|_{L_t^2(H^1)} \\
& + C\|\nabla h_n\|_{L_t^\infty(H^{s-1})}\|u_n - u_{n-1}\|_{L_t^2(H^1)} \\
& \leq C\|h_n\|_{L_t^\infty(H^s)}\|u_n - u_{n-1}\|_{L_t^2(H^s)} \\
& \leq CE_2\|u_n - u_{n-1}\|_{L_t^2(H^s)}.
\end{aligned}$$

Clearly, we have

$$(3.29) \quad \|J_2\|_{\tilde{L}_t^2(H^{s-1})} = \|\operatorname{div}(u_n - u_{n-1})\|_{L_t^2(H^{s-1})} \leq C\|u_n - u_{n-1}\|_{L_t^2(H^s)}.$$

By Lemma 2.12, we obtain

(3.30)

$$\begin{aligned}
& \|J_3\|_{\tilde{L}_t^2(H^{s-1})} = \|h_n \operatorname{div}(u_n - u_{n-1})\|_{\tilde{L}_t^2(H^{s-1})} \\
& \leq C\|h_n\|_{L_t^\infty(L^\infty)}\|\operatorname{div}(u_n - u_{n-1})\|_{\tilde{L}_t^2(H^{s-1})} + C\|\operatorname{div}(u_n - u_{n-1})\|_{\tilde{L}_t^2(B_{\infty,\infty}^{s-2})}\|h_n\|_{L_t^\infty(H^1)} \\
& + C\|h_n\|_{L_t^\infty(H^1)}\|\operatorname{div}(u_n - u_{n-1})\|_{\tilde{L}_t^2(H^{s-1})} \\
& \leq C\|h_n\|_{L_t^\infty(H^s)}\|u_n - u_{n-1}\|_{\tilde{L}_t^2(H^s)} \\
& \leq CE_2\|u_n - u_{n-1}\|_{\tilde{L}_t^2(H^s)}.
\end{aligned}$$

Similarly to J_3 , by Lemma 2.12, we have

(3.31)

$$\begin{aligned}
& \|J_4\|_{\tilde{L}_t^2(H^{s-1})} = \|(h_n - h_{n-1}) \operatorname{div} u_{n-1}\|_{\tilde{L}_t^2(H^{s-1})} \\
& \leq C\|h_n - h_{n-1}\|_{L_t^\infty(B_{\infty,\infty}^{s-2})}\|\operatorname{div} u_{n-1}\|_{L_t^2(H^1)} + \|\operatorname{div} u_{n-1}\|_{L_t^2(L^\infty)}\|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
& + C\|\operatorname{div} u_{n-1}\|_{L_t^2(H^1)}\|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
& \leq C\|u_{n-1}\|_{L_t^2(H^{s+1})}\|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})} \\
& \leq CE_1\|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})}.
\end{aligned}$$

Thus we have

(3.32)

$$\begin{aligned} \|h_{n+1} - h_n\|_{\tilde{L}_t^\infty(H^{s-1})} &\leq 2(\|\Delta_{n+1}h_0\|_{H^{s-1}} + T_1^{\frac{1}{2}}\|\sum_{j=1}^4 G_j\|_{\tilde{L}_t^2(H^{s-1})}) \\ &\leq 2\|\Delta_{n+1}h_0\|_{H^{s-1}} + CT_1^{\frac{1}{2}}(1 + E_2)\|u_n - u_{n-1}\|_{L_t^2(H^s)} + CT_1^{\frac{1}{2}}E_1\|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})}. \end{aligned}$$

Similarly as $\|u_{n+1} - u_n\|_{\tilde{L}_t^\infty(H^{s-1})}$, for all $\rho_j \leq 2$, we have

(3.33)

$$\begin{aligned} \|u_{n+1} - u_n\|_{L_t^2(H^s)} &\leq \nu^{-\frac{1}{2}}C_0e^{C_0\int_0^t\|u_n\|_{H^{s+1}}dt'}\left((1+\nu t)^{\frac{1}{2}}\|\Delta_{n+1}u_0\|_{H^{s-1}}\right. \\ &\quad \left. + \sum_{j=1}^5(1+\nu t)^{1+\frac{1}{2}-\frac{1}{\rho_j}}\nu^{\frac{1}{\rho_j}-1}\|F_j\|_{\tilde{L}_t^{\rho_j}(H^{s-3+\frac{2}{\rho_j}})})\right) \\ &\leq 4\nu^{-1}C_0\|\Delta_{n+1}u_0\|_{H^{s-1}} + 4\nu^{-1}\sum_{j=1}^5\|F_j\|_{\tilde{L}_t^{\rho_j}(H^{s-3+\frac{2}{\rho_j}})} \\ &\leq 4\nu^{-1}C_0\|\Delta_{n+1}u_0\|_{H^{s-1}} + CT_1^{\frac{s-1}{4}}E_1\|u_n - u_{n-1}\|_{L_t^\infty(H^{s-1})} + CT_1^{\frac{s-1}{4}}E_2\|u_n - u_{n-1}\|_{L_t^2(H^s)} \\ &\quad + CT_1^{\frac{s-1}{4}}(1 + E_1 + E_1E_2 + E_1E_2^2)\|h_n - h_{n-1}\|_{L_t^\infty(H^{s-1})}. \end{aligned}$$

We also have

$$\begin{aligned} (3.34) \quad \|\Delta_{n+1}u_0\|_{H^{s-1}} &\leq 2^{-(n+1)}\|\Delta_{n+1}u_0\|_{H^s} \leq 2^{-(n+1)}\|u_0\|_{H^s} \leq 2^{-(n+1)}E_1, \\ \|\Delta_{n+1}h_0\|_{H^{s-1}} &\leq 2^{-(n+1)}\|\Delta_{n+1}h_0\|_{H^s} \leq 2^{-(n+1)}\|h_0\|_{H^s} \leq 2^{-(n+1)}E_2. \end{aligned}$$

We will complete this proof by induction. By Proposition 3.1, we have

$$\begin{aligned} \|u_2 - u_1\|_{\tilde{L}_{T_1}^\infty(H^{s-1})} &\leq \|u_1\|_{\tilde{L}_{T_1}^\infty(H^{s-1})} + \|u_2\|_{\tilde{L}_{T_1}^\infty(H^{s-1})} \\ &\leq \|u_1\|_{\tilde{L}_{T_1}^\infty(H^s)} + \|u_2\|_{\tilde{L}_{T_1}^\infty(H^s)} \\ &\leq 2E_1. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \|u_2 - u_1\|_{\tilde{L}_{T_1}^2(H^s)} &\leq 2E_1, \\ \|h_2 - h_1\|_{\tilde{L}_{T_1}^\infty(H^{s-1})} &\leq 2E_2. \end{aligned}$$

We assume that

$$\|u_n - u_{n-1}\|_{\tilde{L}_{T_1}^\infty(H^{s-1})} \leq 8 \times 2^{-n} E_1,$$

$$\|u_n - u_{n-1}\|_{\tilde{L}_{T_1}^2(H^s)} \leq 8 \times 2^{-n} E_1,$$

$$\|h_n - h_{n-1}\|_{\tilde{L}_{T_1}^\infty(H^{s-1})} \leq 8 \times 2^{-n} E_2.$$

Thus take a suitable T_1 satisfying

$$(3.35) \quad \begin{cases} 8CT_1^{\frac{s-1}{4}} E_1 \leq 1, & 8CT_1^{\frac{s-1}{4}} E_2 \leq 1, \\ 8CT_1^{\frac{s-1}{4}} (1 + E_1 + E_1 E_2 + E_1 E_2^2) E_2 E_1^{-1} \leq 1, \\ 8CT_1^{\frac{1}{2}} E_1 \leq 1, & 8CT_1^{\frac{1}{2}} E_1 (1 + E_2) E_2^{-1} \leq 1. \end{cases}$$

Where C in subsection 3.2 only depends on C_0, ν as well. Thus using (3.26), (3.32), (3.33), we can obtain that

$$\|u_{n+1} - u_n\|_{\tilde{L}_{T_1}^\infty(H^{s-1})} \leq 8 \times 2^{-(n+1)} E_1,$$

$$\|u_{n+1} - u_n\|_{\tilde{L}_{T_1}^2(H^s)} \leq 8 \times 2^{-(n+1)} E_1,$$

$$\|h_{n+1} - h_n\|_{\tilde{L}_{T_1}^\infty(H^{s-1})} \leq 8 \times 2^{-(n+1)} E_2.$$

This completes the proof of Proposition 3.3.

3.3. Existence and uniqueness of local solutions

In this subsection, we investigate the uniqueness of the local solution to the system (2.1). By Proposition 3.3, the approximative sequence (u_n, h_n) of the problem (3.1) is a Cauchy sequence in $\chi([0, T_1], s-1, E_1, E_2)$ with $s \in (1, 2)$. So the limit (u, h) is a solution of the Cauchy problem (2.1). From Proposition 3.1, we obtain that this sequence is bounded in $\chi([0, T], s, E_1, E_2)$, so it's also the Cauchy sequence in $\chi([0, T_1], s', E_1, E_2)$ for all $s' < s$ by interpolation, and the limit is in $\chi([0, T_1], s, E_1, E_2)$. Thus we have proved local existence result in Theorem 1.1.

For the uniqueness result in Theorem 1.1, let (u, h) and (v, g) satisfy the problem (2.1) with the initial data $(u_0, h_0), (v_0, g_0) \in H^s \times H^s$ respectively. Then we have

$$(3.36) \quad \begin{cases} \partial_t(u - v) + u \cdot \nabla(u - v) - \nu \Delta(u - v) = G_1(u, h) - G_1(v, g), \\ \partial_t(h - g) + u \cdot \nabla(h - g) = (u - v) \nabla g + G_2(u, h) - G_2(v, g), \\ (u - v)|_{t=0} = 0, (h - g)|_{t=0} = 0. \end{cases}$$

Using Lemmas 2.14-2.15, similarly as the proof of Proposition 3.3, for all $t \leq T_1$, we can get,

$$(3.37) \quad \begin{aligned} & \|u - v\|_{\tilde{L}^\infty(H^s)} + \|u - v\|_{\tilde{L}^2(H^{s+1})} + \|h - g\|_{\tilde{L}^\infty(H^s)} \\ & \leq C e^{Ct} (\|u_0 - v_0\|_{H^s} + \|h_0 - g_0\|_{H^s}) \\ & \quad + C t^{\alpha_1} \|u - v\|_{\tilde{L}^\infty(H^s)} + C t^{\alpha_2} \|u - v\|_{\tilde{L}^2(H^{s+1})} + C t^{\alpha_3} \|h - g\|_{\tilde{L}^\infty(H^s)}, \end{aligned}$$

where $\alpha_1, \alpha_2, \alpha_3 \in (0, 1)$, and C only depends on C_0, ν, E_1, E_2 . Taking a suitable $T_2 \leq T_1$ gives the uniqueness of Theorem 1.1 in $[0, T_2]$, and thus we get the uniqueness in $[0, T_1]$.

4 Global existence

In [12], Wang and Xu got the global existence of the Cauchy problem (1.1) for small initial data in $H^s \times H^s$ with $s > 2$, since they showed only the existence of the local solutions in $H^s \times H^s$ with $s > 2$. And They also pointed out that (see Remark of Theorem 4.2 in [12]): if one could establish the local solutions in $H^s \times H^s$ with $s > 1$, one can also obtain the global solution with initial data in $H^s \times H^s$ with $s > 1$. Since the proof of global existence is the same as that of the $s > 2$ case [12], here we omit it.

Remark 4.1 Note Wang and Xu in [12] established the local well-posedness of the system (1.1) for any initial data and got the global solutions to the system (1.1) for

small initial data in Sobolev spaces H^s , $s > 2$. Our obtained result in Theorem 1.1 shows that Wang and Xu's conjecture in [12] holds true.

Acknowledgements. This work was partially supported by NNSFC (No.11271382 No. 10971235), RFDP (No. 20120171110014), and the key project of Sun Yat-sen University.

References

- [1] Adams, R. A. and Fournier, J. F., *Sobolev spaces*, Academic Press, New York, (2003).
- [2] Bahouri, H., Chemin, J-Y. and Danchin, R., *Fourier Analysis and Nonlinear Partial Differential Equations*, Springer, Grundlehren der mathematischen Wissenschaften, 343 (2011), 51-164.
- [3] Bresch, D. and Desjardins, B., *Existence of global weak solutions for 2D viscous shallow water equations and convergence to the quasi-geostrophic model*, Commun. Math. Phys, **238:1-2** (2003), 211-233.
- [4] Bresch, D., Desjardins, B. and Metivier, G., *Recent mathematical results and open problem about shallow water equations*, Analysis and simulation of Fluid Dynamics, Advances in Mathematical Fluid Mechanics, 15-31, Birkhäuser Verlag Basel/Switzerland, 2006.
- [5] Bui, A.-T., *Existence and uniqueness of a classical solution of an initial boundary value problem of the theory of shallow waters*, SIAM J. Math. Anal., **12** (1981) 229-241.

- [6] Chen, Q., Miao, C., and Zhang, Z., *Well-posedness for the viscous shallow water equations in critical spaces*, SIAM J. Math. Anal. **40** (2008), 443-474.
- [7] B. Haspot, *Cauchy problem for viscous shallow water equations with a term of capillarity*, Hyperbolic problems: theory, numerics and applications, 625-634, Proc. Sympos. Appl. Math., 67, Part 2, Amer. Math. Soc., Providence, RI, 2009.
- [8] Kloeden, P.-E., *Global existence of classic solution in the dissipative shallow water equations*, SIAM J. Math. Anal., **16** (1985), 301-315.
- [9] Matsumura, A., Nishida, T., *The initial value problem for the equations of motion of viscous and heat-conductive gases*, J. Math. Kyoto Univ., **20** (1980), 67-104.
- [10] Sondbye, L., *Global existence for Dirichlet problem for the viscous shallow water equations*, J. Math. Anal. Appl., **202** (1996), 236-258.
- [11] Sondbye, L., *Global existence for the Cauchy problem for the viscous shallow water equations*, Rocky Mountain J. Math., **28** (1998), 1135-1152.
- [12] Wang, W. and Xu, C.-J., *The Cauchy problem for viscous shallow water equations*, Rev. Mat. Iberoamericana, **21:1** (2005), 1-24.